	_										
Reg. No. :											



.()

## Question Paper Code: 91783

B.E./B.Tech. DEGREE EXAMINATIONS, NOVEMBER/DECEMBER 2019
Third Semester

Civil Engineering

MA 6351 – TRANSFORMS AND PARTIAL DIFFERENTIAL EQUATIONS (Common to All Branches)

(Regulations 2013)

Time: Three Hours

Maximum: 100 Marks

Answer ALL questions.

PART - A

 $(10\times2=20 \text{ Marks})$ 

- 1. Form the partial differential equation by eliminating the arbitrary constants  $\hat{a}$  and b from  $z = (x a)^2 + (y b)^2 + 1$ .
- 2. Find the complete integral of p + q = x + y.
- 3. State the Dirichlet's conditions for a function f(x) to be expanded as a Fourier series.
- 4. Expand f(x) = 1, in  $(0, \pi)$  as a half-range sine series.
- 5. Write all possible solutions of one dimensional heat equation  $\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$ .
- 6. Using the method of separation of variables, solve  $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} + u$  where  $u(x, 0) = 6e^{-3x}$ .
- 7. If the Fourier transform of f(x) is  $\Im(f(x)) = F(s)$ , then show that  $\Im(f(x-a)) = e^{ias} F(s)$ .
- 8. Find the Fourier sine transform of 1/x.
- 9. Find the Z transform of  $\frac{1}{n+1}$
- 10. State the final value theorem of Z transforms.

(8)

( )

PART - B

(5×16=80 Marks)

- 11. a) i) Find the general solution of  $(z^2 2yz y^2)p + (xy + zx)q = xy zx$ . (8)
  - ii) Find the general solution of  $(D^2 + 2DD' + D'^2)$   $z = x^2 y + e^{x-y}$ . (8)

(OR)

- b) i) Find the general solution of  $z = px + qy + p^2 + pq + q^2$ . (8)
  - ii) Find the general solution of  $(D^2 3DD' + 2D'^2 + 2D 2D')$  z = sin (2x + y). (8)
- 12. a) i) Find the Fourier series expansion of the following periodic function  $f(x) = \begin{cases} 2+x-2 \le x \le 0 \\ 2-x & 0 < x \le 2 \end{cases}$  of period 4 Hence deduce that  $\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{2}.$ 
  - ii) Find the complex form of Fourier series of  $f(x) = e^{ax}$  in the interval  $(-\pi, \pi)$  where a is a real constant. Hence, deduce that  $\sum_{n=-\infty}^{\infty} \frac{(-1)^n}{a^2 + n^2} = \frac{\pi}{a \sinh a\pi}.$  (8)
  - b) i) Find the half range cosine series of  $f(x) = (\pi x)^2$ ,  $0 < x < \pi$ . Hence find the sum of the series  $\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots$  (8)
    - ii) Determine the first two harmonics of Fourier series for the following data.  $x : 0 \quad \frac{\pi}{3} \quad \frac{2\pi}{3} \quad \pi \quad \frac{4\pi}{3} \quad \frac{5\pi}{3}$ (8)

f(x): 1.98 1.30 1.05 1.30 -0.88 -0.25

- 13. a) A tightly stretched string of length 'l' with fixed end points is initially at rest in its equilibrium position. If it is set vibrating by giving each point a velocity  $y_l(x,0) = v_0 \sin\left(\frac{3\pi x}{l}\right) \cos\left(\frac{\pi x}{l}\right), \text{ where } 0 < x < l. \text{ Find the displacement of the string at a point, at a distance x from one end at any instant 't'.} \tag{16}$ 
  - b) A square plate is bounded by the lines x = 0, x = 20, y = 0, y = 20. Its faces are insulated. The temperature along the upper horizontal edge is given by u(x, 20) = x (20 x), 0 < x < 20, while the other three edges are kept at 0°C. Find the steady state temperature distribution u(x, y) in the plate. (16)</li>

14. a) i) Find the Fourier Transform of f(x) if  $f(x) = \begin{cases} 1 - |x|, |x| < 1 \\ 0, |x| > 1 \end{cases}$  and hence evaluate the integral  $\int_{0}^{\infty} \left(\frac{\sin t}{t}\right)^{4} dt$ . (10)

ii) State and prove convolution theorem for Fourier transforms. (6)
(OR)

b) i) Evaluate  $\int_{0}^{\infty} \frac{dx}{(x^2 + a^2)(x^2 + b^2)}$  using transforms. (6)

ii) Find the Fourier cosine transform of  $f(x) = e^{-a^2x^2}$  and hence

find  $F_s\left[xe^{-a^2x^2}\right]$ . (10)

15. a) i) Find  $Z(r^n \cos n\theta)$  and  $Z^{-1} \left[ \left( 1 - az^{-1} \right)^{-2} \right]$ . (8)

ii) Using convolution theorem, find  $Z^{-1}\left[\frac{z^2}{(z-1/2)(z-1/4)}\right]$ . (8)

(OR)

- b) i) Using Z-transform, solve the difference equation x(n + 2) 3x(n + 1) + 2x(n) = 0 given that x(0) = 0, x(1) = 1.
  - ii) Using residue method, find  $Z^{-1} \left[ \frac{z}{z^2 2z + 2} \right]$ . (8)